





WO9709049A2: METHOD FOR EFFECTING VASODILATION WITH (1,5-INTER)ARYL PROSTAGLANDIN DERIVATIVES

View Images (31 pages) | View Cart

Premium Data 1: PDF (~2950 KB) | TIFF (~2330 KB) | Fax | More choices...



Profity Numbers

Designated

CHEN, June., 28771 Via Los Arboles, San Juan Capistrano, CA 92675, United States of

BURK, Robert, M., 1337 Cerritos Drive, Laguna Beach, CA 92651, United States of

WOODWARD, David, F., 23152 Tulip, Lake Forest, CA 92630, United States of America

ALLERGAN, 8301 Mars Drive, Waco, TX 76712, United States of America

March 13, 1997 / Aug. 28, 1996

WO1996US0013889

A61K 031/557; C07C 405/00;

Sept. 1, 1995 US1995008522775

AU, CA, JP, European patent: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE



The present invention provides a method of effecting vasodilation, comprising: administering to a warm blooded animal in need of such treatment, an effective amount of (1,5-inter)aryl prostaglandin derivative represented by formula (I), wherein n is 0 or an integer of from 1 to 6, R is selected from the group of radicals represented by the formulae CO2R', CONR'2, CH2OR' and SO2NR'2', wherein R' is hydrogen or a lower alkyl radical having fromone to six carbon atoms; R" is hydrogen or an acyl radical having the formula: (CO)R" wherein R" is a saturated or unsarrurated acyclic hydrocarbon radical having from 1 to about 10 carbon atoms, or -(CH2)mR"" wherein m is 0 or an integer of from 1 to 6 and R"" is an aliphatic ring having from 3 to 7 carbon atoms or an aryl group, e.g. phenyl, or a heteroaryl group, e.g. thienyl, furanyl or pyridyl, and preferably R" is a lower alkyl group having from 1 to 6 carbon atoms; the hatched triangular segments represent alpha oriente bonds, the solid triangular segments represent beta oriented bonds and the wavy segmen represent bonds that may be in either thecis or trans orientation. More preferably said (1,5 inter)aryl prostaglandin derivative is a compound represented by formula (II). Most preferably, said (1,5-inter)aryl prostaglandin derivative is a compound of formula (II) wherein R is CO2R', R' is hydrogen and n is 0. [Show "fr" Abstract]

